



## **Clinical Opinion**

## Ultrasound guided intra-articular botulinum toxin injection for treatment of the osteoarthritic knee pain: expert opinion.

Allam Abdallah El-Sayed <sup>1,2\*</sup>, Galluccio Felice <sup>3</sup>.

 Department of physical medicine Rheumatology and rehabilitation Tanta University Hospital Egypt
Egypt
Department of rheumatology, department of clinical and experimental medicine AOU Careggi Florence Italy
\*Corresponding author Correspondence to: allmovish@gmail.com. Publication data: Submitted: October 23,2019 Accepted: February 27,2020 Online: March 15.2020

This article was subject to full peer-review

This is an open access article distributed under the terms of the Creative Commons Attribution Non-Commercial License 4.0 (CCBY-NC) allowing to share

Share: copy and redistribute the material in any

Adapt: remix, transform, and build upon the licensed

the work provided must be properly cited and cannot

and adapt.

material

medium or format.

be used for commercial purpose.

Osteoarthritis is the most common form of arthritis and cause of joint pain and disability [1]. The prevalence, burden and socioeconomic impact of osteoarthritis of the knee is considerable and growing [2]. Knee pain is the most cardinal symptom of the osteoarthritic knee and is associated with a high degree of impaired function and daily activities [3]. Intra-articular injection provides a safe and cost effective control of knee pain with improvement of function and activities of daily living [4]. Ultrasound guidance improves the accuracy of the intra-articular injection in knee joint and clinical outcomes and lowers health care costs [5]. Botulinum toxin targets many neuropeptides leading to reduction of pain, peripheral sensitization and in turn reduction of the central sensitization. Also, botulinum toxin plays an anti-inflammatory role by reduction of the neurogenic inflammation [6]. The following table summarizes how to use botulinum toxin in clinical practice according to authors experience. However, further large-scale multicenter randomized controlled trials are needed to confirm effectiveness and determine the exact position of this intervention in the therapeutic armamentarium for the osteoarthritic knee pain.

DOSE PER KNEE	COMPLICATIONS AND CONTRA - INDICATIONS	FREQUENCY OF INJECTIONS	DURATION OF PAIN RELIEF	FAILURE OF RESPONSE	COMBINATIONS
Botox → 50 IU Dysport → 100 IU	Similar to general complications and contra-indications as a regard local injection	Can be repeated after 6 months	* 6 - 12 months after 1st injection * 12 - 18 months after 2nd injection	After second injection due to auto-antibodies formation is rare.	Can be combined with hyaluronic acid and platele rich plasma
Botox $\rightarrow$ 100 IU Dysport $\rightarrow$ 200 IU		Onset of action			
Botox → 200 IU Dysport → 400 IU		Delayed. Patients might start to feel the improvement of the pain after 2 weeks of the injection.			

-----, -

## References:

[1] Damen J, van Rijn RM, Emans PJ, Hilberdink WKHA, Wesseling J, Oei EHG, et al. Prevalence and development of hip and knee osteoarthritis according to American College of Rheumatology criteria in the CHECK cohort. Arthritis Res Ther. 2019; 21:4.

[2] Emery CA, Whittaker JL, Mahmoudian A, Lohmander LS, Roos EM, Bennell KL, et al. Establishing outcome measures in early knee osteoarthritis. Nat Rev Rheumatol. 2019; 15:438-48.

[3] Conoghan PG, Cook AD, Hamilton JA, Tak PP. Therapeutic options for targeting inflammatory osteoarthritis pain. Nat Rev Rheumatol. 2019; 15:355-63.

[4] Tapasvi S, Mohanty SS, Vedavyasa Acharya KK, Bhattacharya K, et al. Viscosupplementation for Management of Knee Osteoarthritis from an Indian Perspective: An Expert Consensus Report. Pain Ther. 2019; 8:217-231.

[5] Berkoff DJ, Miller LE, Block JE. Clinical utility of ultrasound guidance for intra-articular knee injections: a review. Clin Interv Aging. 2012 ;7 :89-95.

[6] Khenioui H, Houvenagel E, Catanzariti JF, Guyot MA, Agnani O, Donze C. Usefulness of intra-articular botulinum toxin injections. A systematic review. Joint Bone Spine. 2016; 83:149-54.

Citation: Allam AE, Galluccio F. Ultrasound guided intra-articular botulinum toxin injection for treatment of the osteoarthritic knee pain; expert opinion. Jr.med.res. 2020; 3(1):25. Allam © All rights are reserved. <u>https://doi.org/10.32512/jmr.3.1.2020/25</u> Submit your manuscript: <u>www.jmedicalresearch.com</u>